




## Separation of Neonicotinoids on HALO® C18, 2.7 µm

Application Note 92-PS



### PEAK IDENTITIES:

1. Nitenpyram
2. Thiamethoxam
3. Clothianidin
4. Imidacloprid
5. Acetamiprid
6. Thiachloprid

Neonicotinoids are systemic insect neurotoxins that have recently been in the news, since this class of pesticides may have negative effects on bees. This application note shows a rapid separation of six neonicotinoids using a Fused-Core®, 2.7 µm, HALO® C18 column. This superficially porous packing allows high resolution at moderate back pressures.

### TEST CONDITIONS:

**Column:** HALO 90 Å C18, 2.7 µm,  
3.0 x 100 mm

**Part Number:** 92813-602

**Mobile Phase:** 70/30 - A/B

A: 0.1% formic acid in water

B: Acetonitrile

**Flow Rate:** 0.8 mL/min

**Pressure:** 252 bar

**Temperature:** 35 °C

**Detection:** UV 254 nm, VWD

**Injection Volume:** 2.0 µL

**Sample Solvent:** 50/50 water/acetonitrile


**Response Time:** 0.02 sec

**Flow Cell:** 2.5 µL semi-micro


**LC System:** Shimadzu Prominence UFLC XR

**Extra Column Volume:** ~14 µL


### STRUCTURES:




Nitenpyram




Imidacloprid




Thiamethoxam



Acetamiprid



Clothianidin



Thiachloprid

